

The fractional volatility model: No-arbitrage, leverage and completeness

R. Vilela Mendes^{*†}, M. J. Oliveira^{*‡} and A.M. Rodrigues^{*}

Abstract

Based on a criterion of mathematical simplicity and consistency with empirical market data, a stochastic volatility model has been obtained with the volatility process driven by fractional noise. Depending on whether the stochasticity generators of log-price and volatility are independent or are the same, two versions of the model are obtained with different leverage behavior. Here, the no-arbitrage and completeness properties of the models are studied.

Keywords: Fractional noise, Arbitrage, Incomplete market

1 Introduction

In liquid markets the autocorrelation of price changes decays to negligible values in a few minutes, consistent with the absence of long term statistical arbitrage. Because innovations of a martingale are uncorrelated, there is a strong suggestion that it is a process of this type that should be used to model the stochastic part of the returns process. As a consequence, classical Mathematical Finance has, for a long time, been based on the assumption that the price process of market securities may be approximated by geometric Brownian motion

$$dS_t = \mu S_t dt + \sigma S_t dB(t) \quad (1)$$

^{*}Centro de Matemática e Aplicações Fundamentais, Universidade de Lisboa, Av. Prof. Gama Pinto 2, P1649-003 Lisboa

[†]IPFN, EURATOM/IST Association, e-mail: rvilela.mendes@gmail.com

[‡]Universidade Aberta, oliveira@cii.fc.ul.pt

Geometric Brownian motion (GBM) models the absence of linear correlations, but otherwise has some serious shortcomings. It does not reproduce the empirical leptokurtosis nor does it explain why nonlinear functions of the returns exhibit significant positive autocorrelation. For example, there is volatility clustering, with large returns expected to be followed by large returns and small returns by small returns (of either sign). This, together with the fact that autocorrelations of volatility measures decline very slowly [1], [2], [3] has the clear implication that long memory effects should somehow be represented in the process and this is not included in the geometric Brownian motion hypothesis. The existence of an essential memory component is also clear from the failure of reconstruction of a Gibbs measure and the need to use chains with complete connections in the phenomenological reconstruction of the market process [4].

As pointed out by Engle [5], when the future is uncertain investors are less likely to invest. Therefore uncertainty (volatility) would have to be changing over time. The conclusion is that a dynamical model for volatility is needed and σ in Eq.(1), rather than being a constant, becomes itself a process. This idea led to many deterministic and stochastic models for the volatility ([6], [7] and references therein).

The stochastic volatility models that were proposed described some partial features of the market data. For example leptokurtosis is easy to fit but the long memory effects are much harder. On the other hand, and in contrast with GBM, some of the phenomenological fittings of historical volatility lack the kind of nice mathematical properties needed to develop the tools of mathematical finance. In an attempt to obtain a model that is both consistent with the data and mathematically sound, a new approach was developed in [8]. Starting only with some criteria of mathematical simplicity, the basic idea was to let the data itself tell us what the processes should be.

The basic hypothesis for the model construction were:

(H1) The log-price process $\log S_t$ belongs to a probability product space $(\Omega_1 \times \Omega_2, P_1 \times P_2)$ of which the (Ω_1, P_1) is the Wiener space and the second one, (Ω_2, P_2) , is a probability space to be reconstructed from the data. Denote by $\omega_1 \in \Omega_1$ and $\omega_2 \in \Omega_2$ the elements (sample paths) in Ω_1 and Ω_2 and by $\mathcal{F}_{1,t}$ and $\mathcal{F}_{2,t}$ the σ -algebras in Ω_1 and Ω_2 generated by the processes up to t . Then, a particular realization of the log-price process is denoted

$$\log S_t(\omega_1, \omega_2)$$

This first hypothesis is really not limitative. Even if none of the non-trivial

stochastic features of the log-price were to be captured by Brownian motion, that would simply mean that S_t was a trivial function in Ω_1 .

(H2) The second hypothesis is stronger, although natural. It is assumed that for each fixed ω_2 , $\log S_t(\cdot, \omega_2)$ is a square integrable random variable in Ω_1 .

These principles and a careful analysis of the market data led, in an essentially unique way¹, to the following model:

$$dS_t = \mu_t S_t dt + \sigma_t S_t dB(t) \quad (2)$$

$$\log \sigma_t = \beta + \frac{k}{\delta} \{B_H(t) - B_H(t - \delta)\} \quad (3)$$

the data suggesting values of H in the range $0.8 - 0.9$. In this coupled stochastic system, in addition to a mean value, volatility is driven by fractional noise. Notice that this empirically based model is different from the usual stochastic volatility models which assume the volatility to follow an arithmetic or geometric Brownian process. Also in Comte and Renault [9] and Hu [10], it is fractional Brownian motion that drives the volatility, not its derivative (fractional noise). δ is the observation scale of the process. In the $\delta \rightarrow 0$ limit the driving process would be a distribution-valued process.

The equation (3) leads to

$$\sigma_t = \theta e^{\frac{k}{\delta} \{B_H(t) - B_H(t - \delta)\} - \frac{1}{2} \left(\frac{k}{\delta}\right)^2 \delta^{2H}} \quad (4)$$

with $E[\sigma_t] = \theta > 0$.

The model has been shown [8] to describe well the statistics of price returns for a large δ -range and a new option pricing formula, with "smile" deviations from Black-Scholes, was also obtained. An agent-based interpretation [11] also led to the conclusion that the statistics generated by the model was consistent with the limit order book price setting mechanism.

To our surprise, this data-reconstructed model was met with some hostility by the Mathematical Finance community. Perhaps in part because much of the nice results in this field are based on simple GBM and also because fractional Brownian has been associated to arbitrage. In fact, in the past, several authors tried to describe the long memory effect by replacing in the price process Brownian motion by fractional Brownian motion with $H > 1/2$.

¹Essentially unique in the sense that the empirically reconstructed volatility process is the simplest one, consistent with the scaling properties of the data.

However it was soon realized [12], [13], [14], [15] that this replacement implied the existence of arbitrage. These results might be avoided either by restricting the class of trading strategies [16], introducing transaction costs [17] or replacing pathwise integration by a different type of integration [18] [19]. However this is not free of problems because the Skorohod integral approach requires the use of a Wick product either on the portfolio or on the self-financing condition, leading to unreasonable situations from the economic point of view (for example positive portfolio with negative Wick value, etc.) [20].

The fractional volatility model in Eqs.(2-3) is not affected by these considerations, because it is the volatility process that is driven by fractional noise, not the price process. In fact a no-arbitrage result may be proven. This is no surprise because our requirement (H2) that, for each sample path $\omega_2 \in \Omega_2$, $\log S_t(\cdot, \omega_2)$ is a square integrable random variable in Ω_1 already implies that $\int \sigma_t dB_t$ is a martingale. The square integrability is also essential to guarantee the possibility of reconstruction of the σ process from the data, because it implies [21]

$$\frac{dS_t}{S_t}(\cdot, \omega_2) = \mu_t(\cdot, \omega_2) dt + \sigma_t(\cdot, \omega_2) dB_t \quad (5)$$

Our aim in this paper is to give a solid mathematical construction of the fractional volatility model, discussing existence questions, arbitrage and market completeness.

2 No-arbitrage and incompleteness

Let $(\Omega_1, \mathcal{F}_1, P_1)$ be the complete filtered Wiener probability space, carrying a standard Brownian motion $B = (B_t)_{0 \leq t < \infty}$ and a filtration $\mathbb{F}_1 = (\mathcal{F}_{1,t})_{0 \leq t < \infty}$. Let also $(\Omega_2, \mathcal{F}_2, P_2)$ be another probability space associated to a fractional Brownian motion B_H with Hurst parameter $H \in (0, 1)$ and a filtration $\mathbb{F}_2 = (\mathcal{F}_{2,t})_{0 \leq t < \infty}$ generated by B_H .

Let us now embed these two probability spaces in a product space $(\overline{\Omega}, \overline{\mathcal{F}}, \overline{P})$, where $\overline{\Omega}$ is the Cartesian product $\Omega_1 \times \Omega_2$ and \overline{P} is the product measure $P_1 \otimes P_2$. We also introduce π_1 and π_2 , the projections of $\overline{\Omega}$ onto Ω_1 and Ω_2 , as well as the σ -algebra \mathcal{N} generated by all null sets from the product σ -algebra $\mathcal{F}_1 \otimes \mathcal{F}_2$, that is,

$$\mathcal{N} = \sigma(\{F \subseteq \Omega_1 \times \Omega_2 | \exists G \in \mathcal{F}_1 \otimes \mathcal{F}_2 \text{ such that } F \subseteq G \text{ and } (P_1 \otimes P_2)(G) = 0\}).$$

Moreover, we let $\overline{\mathcal{F}} = (\mathcal{F}_1 \otimes \mathcal{F}_2) \vee \mathcal{N}$, the σ -algebra generated by the union of the σ -algebras $\mathcal{F}_1 \otimes \mathcal{F}_2$ and \mathcal{N} . Then $\overline{\mathbb{F}} = (\overline{\mathcal{F}}_t)_{0 \leq t < \infty}$ is the filtration for $\overline{\mathcal{F}}_t = (\mathcal{F}_{1,t} \otimes \mathcal{F}_{2,t}) \vee \mathcal{N}$.

Furthermore, we extend B and B_H to $\overline{\mathbb{F}}$ -adapted processes on $(\overline{\Omega}, \overline{\mathcal{F}}, \overline{P})$ by $\overline{B}(\omega_1, \omega_2) = (B \circ \pi_1)(\omega_1, \omega_2)$ and $\overline{B}_H(\omega_1, \omega_2) = (B_H \circ \pi_2)(\omega_1, \omega_2)$ for $(\omega_1, \omega_2) \in \overline{\Omega}$. Then, it is easy to prove that \overline{B} and \overline{B}_H are Brownian and fractional Brownian motions with respect to \overline{P} and are independent. For notational simplicity, hereafter B and B_H will stand for \overline{B} and \overline{B}_H .

We now consider a market with a risk-free asset with dynamics given by

$$dA_t = rA_t dt \quad A_0 = 1 \quad (6)$$

with $r > 0$ constant and a risky asset with price process S_t given by Eqs.(2)-(3), with μ_t a $\overline{\mathbb{F}}$ -adapted process with continuous paths, k the volatility intensity parameter and δ the observation time scale of the process.

Eq.(3) leads to Eq.(4)

$$\sigma_t = \theta e^{\frac{k}{\delta}\{B_H(t) - B_H(t-\delta)\} - \frac{1}{2}\left(\frac{k}{\delta}\right)^2 \delta^{2H}}$$

with $\theta = \mathbb{E}_{\overline{P}}[\sigma_t]$, hence σ_t is a measurable and an $\overline{\mathbb{F}}$ -adapted process satisfying for all $0 \leq t < \infty$

$$\begin{aligned} \mathbb{E}_{\overline{P}} \left[\int_0^t \sigma_s^2 ds \right] &= \int_0^t \theta^2 e^{-\left(\frac{k}{\delta}\right)^2 \delta^{2H}} \mathbb{E}_{\overline{P}} \left[e^{\frac{2k}{\delta}\{B_H(s) - B_H(s-\delta)\}} \right] ds \\ &= \theta^2 \exp \left\{ \left(\frac{k}{\delta} \right)^2 \delta^{2H} \right\} t < \infty \end{aligned}$$

using Fubini's theorem and the moment generating function of the Gaussian random variable $B_H(s) - B_H(s - \delta)$.

Moreover $\int_0^t |\mu_s| ds$ being finite \overline{P} -almost surely for $0 \leq t < \infty$, an application of Itô's formula yields

$$S_t = S_0 \exp \left\{ \int_0^t \left(\mu_s - \frac{1}{2} \sigma_s^2 \right) ds + \int_0^t \sigma_s dB_s \right\}$$

Lemma 2.1. *Consider the measurable, $\overline{\mathbb{F}}$ -adapted process defined by*

$$\gamma_t = \frac{r - \mu_t}{\sigma_t}, \quad 0 \leq t < \infty \quad (7)$$

with $\mu \in L^\infty([0, T] \times \overline{\Omega})$ and denote by $\eta = (\eta_t)_{0 \leq t < \infty}$ the stochastic exponential of $\left(\int_0^t \gamma_s dB_s\right)_{0 \leq t < \infty}$, that is,

$$\eta_t = \exp \left\{ \int_0^t \gamma_s dB_s - \frac{1}{2} \int_0^t \gamma_s^2 ds \right\}, \quad 0 \leq t < \infty \quad (8)$$

Then,

$$\mathbb{E}_{\overline{P}} \left[\frac{1}{2} \int_0^T \gamma_s^2 ds \right] < \infty, \quad 0 \leq T < \infty \quad (9)$$

Proof: We make of use the fact that the sample paths of the fractional Brownian motion B_H are Hölder continuous of any order $\alpha \geq 0$ strictly less than H , that is, there is $C_\alpha > 0$ such that, \overline{P} -almost surely, $|B_H(t) - B_H(s)| \leq C_\alpha |t - s|^\alpha$, for every $t, s \in [0, \infty)$. Then, for all $0 \leq T < \infty$

$$\begin{aligned} & \mathbb{E}_{\overline{P}} \left[\frac{1}{2} \int_0^T \gamma_s^2 ds \right] \\ & \leq \mathbb{E}_{\overline{P}} \left[\exp \left\{ \frac{e^{k^2 \delta^{2H-2}}}{2\theta^2} \int_0^T (r + |\mu_s|)^2 e^{-2\frac{k}{\delta}(B(s)-B(s-\delta))} ds \right\} \right] \\ & \leq \mathbb{E}_{\overline{P}} \left[\exp \left\{ \frac{(r + \|\mu_s\|_\infty)^2}{2\theta^2} e^{k^2 \delta^{2H-2}} \int_0^T e^{-2\frac{k}{\delta}(B(s)-B(s-\delta))} ds \right\} \right] \\ & \leq \exp \left\{ \frac{T(r + \|\mu_s\|_\infty)^2}{2\theta^2} e^{k^2 \delta^{2H-2} - 2C_\alpha k \delta^{\alpha-1}} \right\} < \infty \end{aligned}$$

■

Additionally, we assume that investors are allowed to trade only up to some fixed finite planning horizon $T > 0$.

Proposition 2.2. *The market defined by (2), (3) and (6) is free of arbitrage*

Proof: Because, by the lemma above the process γ in (7) satisfies the Novikov condition (9), the nonnegative continuous supermartingale η in (8) is a true \overline{P} -martingale. Hence we can define for each $0 \leq T < \infty$ a new probability measure Q_T on $\overline{\mathcal{F}}_T$ by

$$\frac{dQ_T}{d\overline{P}} = \eta_T, \quad \overline{P} - a.s. \quad (10)$$

Then, by the Cameron-Martin-Girsanov theorem, for each fixed $T \in [0, \infty)$, the process

$$B_t^* = B_t - \int_0^t \frac{r - \mu_s}{\sigma_s} ds \quad 0 \leq t \leq T \quad (11)$$

is a Brownian motion on the probability space $(\bar{\Omega}, \bar{\mathcal{F}}_T, Q_T)$.

Consider now the discounted price process

$$Z_t = \frac{S_t}{A_t} \quad 0 \leq t \leq T$$

Under the new probability measure Q_T , equivalent to \bar{P} on $\bar{\mathcal{F}}_T$, its dynamics is given by

$$Z_t = Z_0 + \int_0^t \sigma_s Z_s dB_s^* \quad (12)$$

and is a martingale in the probability space $(\bar{\Omega}, \bar{\mathcal{F}}_T, Q_T)$ with respect to the filtration $(\bar{\mathcal{F}}_t)_{0 \leq t \leq T}$. By the fundamental theorem of asset pricing, the existence of an equivalent martingale measure for Z_t implies that there are no arbitrages, that is, $\mathbb{E}_{Q_T} [Z_t | \bar{\mathcal{F}}_s] = Z_s$ for $0 \leq s < t \leq T$. ■

Another important concept is market completeness. We note that, in this financial model, trading takes place only in the stock and in the money market and, as a consequence, volatility risk cannot be hedged. Hence, since there are more sources of risk than tradable assets, in this model, the market is incomplete, as proved in the next proposition.

Proposition 2.3. *The market defined by (2), (3) and (6) is incomplete*

Proof: Here we use an integral representation for the fractional Brownian motion [22], [23]

$$B_H(t) = \int_0^t K_H(t, s) dW_s \quad (13)$$

W_t being a Brownian motion with respect to \bar{P} , independent from B_t and K is the square integrable kernel

$$K_H(t, s) = C_H s^{\frac{1}{2}-H} \int_s^t (u-s)^{H-\frac{3}{2}} u^{H-\frac{1}{2}} du, \quad s < t$$

$(H > 1/2)$. Then the process

$$\eta'_t = \exp \left(W_t - \frac{1}{2}t \right) \quad (14)$$

is a square-integrable \bar{P} -martingale. Then, defining a standard bi-dimensional Brownian motion,

$$W_t^* = (B_t, W_t)$$

the process $\eta_t^* = \eta_t \eta_t'$

$$\eta_t^* = \exp \left\{ \int_0^t \Gamma_s \bullet dW_t^* - \frac{1}{2} \int_0^t \|\Gamma_s\|^2 ds \right\}$$

where, by lemma 2.1, $\Gamma = (\gamma, 1)$ satisfies the Novikov condition, is also a \overline{P} -martingale. Then, by the Cameron-Martin-Girsanov theorem, the process

$$\widetilde{W}_t^* = \left(\widetilde{W}_t^{*(1)}, \widetilde{W}_t^{*(2)} \right)$$

defined by

$$\begin{aligned} \widetilde{W}_t^{*(1)} &= B_t - \int_0^t \gamma_s ds \\ \widetilde{W}_t^{*(2)} &= W_t - t \end{aligned}$$

is a bi-dimensional Brownian motion on the probability space $(\overline{\Omega}, \overline{\mathcal{F}}_T, Q_T^*)$, where Q_T^* is the probability measure

$$\frac{dQ_T^*}{d\overline{P}} = \eta_T^* = \eta_T \eta_T' \quad (15)$$

Moreover, the discounted price process Z remains a martingale with respect to the new measure Q_T^* . Q_T^* being an equivalent martingale measure distinct from Q_T , the market is incomplete. ■

As stated above, incompleteness of the market is a reflection of the fact that there are two different sources of risk and only one of the risks is tradable. A choice of measure is how one fixes the volatility risk premium.

3 Leverage and completeness

The following nonlinear correlation of the returns

$$L(\tau) = \langle |r(t+\tau)|^2 r(t) \rangle - \langle |r(t+\tau)|^2 \rangle \langle r(t) \rangle \quad (16)$$

is called *leverage* and the *leverage effect* is the fact that, for $\tau > 0$, $L(\tau)$ starts from a negative value whose modulus decays to zero whereas for $\tau < 0$ it has almost negligible values. In the form of Eqs.(2)(3) the volatility process

σ_t affects the log-price, but is not affected by it. Therefore, in its simplest form the fractional volatility model contains no leverage effect.

Leverage may, however, be implemented in the model in a simple way [24] if one identifies the Brownian processes B_t and W_t in (2) and (13). Identifying the random generator of the log-price process with the stochastic integrator of the volatility, at least a part of the leverage effect is taken into account.

The identification of the two Brownian processes means that now, instead of two, there is only one source of risk. Hence it is probable that in this case completeness of the market might be achieved.

Identifying the random generators, a new fractional volatility model is defined [24]

$$\begin{aligned} dS_t &= \mu_t S_t dt + \sigma_t S_t dW_t \\ \log \sigma_t &= \beta + k' \int_{-\infty}^t (t-u)^{H-\frac{3}{2}} dW_u \end{aligned} \quad (17)$$

Proposition 3.1. *The market defined by (17) and (6) is free of arbitrage and complete.*

Proof: The proof of the first part of the proposition is analogous to that of proposition 2.2. In fact, a similar argument to Lemma 2.2 yields that

$$\eta_t = \exp \left\{ \int_0^t \frac{r - \mu_s}{\sigma_s} dW_s - \frac{1}{2} \int_0^t \left(\frac{r - \mu_s}{\sigma_s} \right)^2 ds \right\}$$

is a P_1 -martingale with respect to $(\mathcal{F}_{1,t})_{0 \leq t < T}$ and the probability measure Q_T , defined by $\frac{dQ_T}{dP_1} = \eta_T$ is an equivalent martingale measure.

Now that we have shown that the set of equivalent local martingale measures for the market is non-empty, let Q^* be an element in this set. Then, recalling that $(\mathcal{F}_{1,t})_{0 \leq t < T}$ is the augmentation of the natural filtration of the Brownian motion W_t , by the Girsanov converse [25] [26] there is a $(\mathcal{F}_{1,t})_{0 \leq t < T}$ progressively measurable \mathbb{R} -valued process ϕ such that the Radon-Nikodym density of Q^* with respect to P_1 equals

$$\frac{dQ^*}{dP_1} = \exp \left\{ \int_0^T \phi_s dW_s - \frac{1}{2} \int_0^T \phi_s^2 ds \right\}$$

Moreover the process W_t^* given by

$$W_t^* = W_t - \int_0^t \phi_s ds$$

is a standard Q^* -Brownian motion and the discounted price process Z satisfies the following stochastic differential equation

$$dZ_t = (\mu_t - r + \sigma_t \phi_t) Z_t dt + \sigma_t Z_t dW_t^*$$

Because Z_t is a Q^* -martingale, then it must be hold $\mu(t, \omega) - r + \sigma(t, \omega) \phi(t, \omega) = 0$ almost everywhere w.r.t. $meas \times P$ in $[0, T] \times \Omega$, $meas$ being the Lebesgue measure on the line. It implies

$$\phi(t, \omega) = \frac{r - \mu(t, \omega)}{\sigma(t, \omega)}$$

a. e. $(t, \omega) \in [0, T] \times \Omega_1$. Hence $Q^* = Q$, that is, Q is the unique equivalent martingale measure. This market model is complete. ■

4 Remarks and conclusions

1) Partially reconstructed from empirical data, the fractional volatility model describes well the statistics of returns. The fact that, once the parameters are adjusted by the data for a particular observation time scale δ , it describes well different time lags is related to the fact that the volatility is driven not by fractional Brownian motion but its increments.

Specific trader strategies and psychology should play a role on market crisis and bubbles. However, the fact that in the fractional volatility model the same set of parameters would describe very different markets [8] seems to imply that the market statistical behavior (in normal days) is more influenced by the nature of the financial institutions (the double auction process) than by the traders strategies [11]. Therefore some kind of universality of the statistical behavior of the bulk data throughout different markets would not be surprising.

The identification of the Brownian process of the log-price with the one that generates the fractional noise driving the volatility, introduces an asymmetric coupling between σ_t and S_t that is also exhibited by the market data.

2) In this paper, mathematical consistency of, both versions, of the fractional volatility model has been established. This and its better consistency with the experimental data, makes it a candidate to replace geometrical Brownian as the standard market model.

References

- [1] Z. Ding, C. W. J. Granger and R. Engle; *A long memory property of stock returns and a new model*, Journal of Empirical Finance 1 (1993) 83–106.
- [2] A. C. Harvey; *Long memory in stochastic volatility*, in *Forecasting Volatility in Financial Markets*. J Knight and S Satchell (Eds), 1998, 307–320. Oxford: Butterworth-Heinemann.
- [3] F. J. Breidt, N. Crato and P. Lima; *The detection and estimation of long memory in stochastic volatility models*, Journal of Econometrics 83 (1998) 325–348.
- [4] R. Vilela Mendes, R. Lima and T. Araújo; *A process-reconstruction analysis of market fluctuations*, International Journal of Theoretical and Applied Finance 5 (2002) 797–821.
- [5] R. F. Engle; *Autoregressive conditional heteroscedasticity with estimates of the variance of United Kingdom inflation*, Econometrica 50 (1982) 987–1007.
- [6] S. J. Taylor; *Modeling stochastic volatility: A review and comparative study*, Mathematical Finance 4 (1994) 183–204.
- [7] R. S. Engle and A. J. Patton; *What good is a volatility model?*, Quantitative Finance 1 (2001) 237–245.
- [8] R. Vilela Mendes and M. J. Oliveira; *A data-reconstructed fractional volatility model*, Economics e-journal, discussion paper 2008-22.
- [9] F. Comte and E. Renault; *Long memory in continuous-time stochastic volatility models*, Mathematical Finance 8 (1998) 291–323.
- [10] Y. Hu; *Option pricing in a market where the volatility is driven by fractional Brownian motion*, in Recent Developments in Mathematical Finance, Ed. J.M. Yong, pp. 49–59, World Scientific, 2002.
- [11] R. Vilela Mendes; *"The fractional volatility model: An agent-based interpretation"*, Physica A: Statistical Mechanics and its Applications, 387 (2008) 3987–3994.

- [12] L. C. G. Rogers; *Arbitrage with fractional Brownian motion*, Mathematical Finance 7 (1997) 95–105.
- [13] A. N. Shiryaev; *On arbitrage and replication for fractal models*, Research Report 20, MaPhySto, Department of Mathematical Sciences, University of Aarhus, Denmark 1998.
- [14] D. M. Salopek; *Tolerance to arbitrage*, Stochastic Processes and Applications 76 (1998) 217–230.
- [15] T. Sottinen; *Fractional Brownian motion, random walks and binary market models*, Finance and Stochastics 5 (2001) 343–355.
- [16] P. Cheridito; *Arbitrage in fractional Brownian motion models*, Finance and Stochastics 7 (2003) 533–553.
- [17] P. Guasoni; *No arbitrage under transaction costs, with fractional Brownian motion and beyond*, Mathematical Finance 16 (2006) 569–582.
- [18] Y. Hu and B. Øksendal; *Fractional white noise calculus and applications to finance*, Infinite Dimensional Analysis Quantum Probability and Related Topics 6 (2003) 1–32.
- [19] R. J. Elliott and J. van der Hoek; *A general fractional white noise theory and applications to finance*, Mathematical Finance 13 (2003) 301–330.
- [20] T. Björk and H. Hult; *A note on Wick products and the fractional Black-Scholes model*, Finance and Stochastics 9 (2005) 197–209.
- [21] D. Nualart; *The Malliavin Calculus and Related Topics*, Springer-Verlag, Berlin 1995.
- [22] L. Decreusefond and A.S. Üstünel; *Stochastic analysis of the fractional Brownian motion*, Potential Analysis 10 (1999) 177–214.
- [23] P. Embrechts and M. Maejima; *Selfsimilar processes*, Princeton University Press, Princeton NJ, 2002.
- [24] R. Vilela Mendes; *A fractional calculus interpretation of the fractional volatility model*, Nonlinear Dynamics 55 (2009) 395–399.

- [25] M. Musiela and M. Rutkowski; *Martingale Methods in Financial Modelling*, vol. 36 of *Stochastic Modelling and Applied Probability*, Springer 2005.
- [26] T. Björk; *Arbitrage Theory in Continuous Time*, Oxford University Press, New York 2009.